

		Dose - Every 4 hours as needed. Do not give more than 5 doses in 24			
Active Ingredient		Infants' Oral Suspension	Children's Oral Suspension	Children's Chewable Tablets	Junior Strength Chewable Tablets
Weight *	Age	Acetaminophen 160 mg (in each 5 mL or 1 tsp)	Acetaminophen 160 mg (in each 5 mL or 1 tsp)	Acetaminophen 80 mg (in each tablet)	Acetaminophen 160 mg (in each tablet)
6-11 lbs	0-3 mos	1.25 mL	-	-	-
12-17 lbs	6-11 mos	2.5 mL	-	-	-
18-23 lbs	12-23 mos	3.75 mL	-	-	-
24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)	2 tablets	-
36-47 lbs	4-5 yrs	-	7.5 mL (1½ tsp)	3 tablets	-
48-59 lbs	6-8 yrs	-	10 mL (2 tsp)	4 tablets	2 tablets
60-71 lbs	9-10 yrs	-	12.5 mL (2½ tsp)	5 tablets	2½ tablets
72-95 lbs	11 yrs	-	15 mL (3 tsp)	6 tablets	3 tablets

mL = milliliter tsp = teaspoonful

*Dosing is based primarily on weight, not age.

		Infants Concentrated Drops for ages 6-23 months	Children's Suspension for ages 2-11 years	Junior Strength Chewables for ages 6-11 years	Junior Strength Tablets for ages 6-11 years
Active Ingredient		ibuprofen 50 mg (NSAID) in each 1.25 mL	ibuprofen 100 mg (NSAID) in each 5 mL	ibuprofen 100 mg (NSAID) in each tablet	ibuprofen 100 mg (NSAID) in each tablet
Indications		Fever reducer; relieves minor aches and pains due to the common cold, flu, headaches, and toothaches	Fever reducer; relieves minor aches and pains due to the common cold, flu, soar throat, headaches, and toothaches	Fever reducer; relieves minor aches and pains due to the common cold, flu, soar throat, headaches, and toothaches	Fever reducer; relieves minor aches and pains due to the common cold, flu, soar throat, headaches, and toothaches
Weight*	Age				
under 6 mos.		Do not use	Do not use	Do not use	Do not use
12-17 pounds	6-11 months	1.25 mL every 6-8 hrs not to exceed 4 doses/24 hrs	Use Infants' Concentrated Drops	Use Infants' Concentrated Drops	Use Infants' Concentrated Drops
18-23 pounds	12-23 months	1.875 mL every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓	↓
24-35 pounds	2-3 years	Use Children's Suspension	1 tsp every 6-8 hrs not to exceed 4 doses/24 hrs	Use Children's Suspension	Use Children's Suspension
36-47 pounds	4-5 years	↓	1½ tsp every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓
48-59 pounds	6-8 years	Use Children's Suspension, or Junior Strength Chewables or Tablets	2 tsp every 6-8 hrs not to exceed 4 doses/24 hrs	2 tablets every 6-8 hrs not to exceed 4 doses/24 hrs	2 tablets every 6-8 hrs not to exceed 4 doses/24 hrs
60-71 pounds	9-10 years	↓	2½ tsp every 6-8 hrs not to exceed 4 doses/24 hrs	2½ tablets every 6-8 hrs not to exceed 4 doses/24 hrs	2 tablets every 6-8 hrs not to exceed 4 doses/24 hrs
72-95 pounds	11 years	↓	3 tsp every 6-8 hrs not to exceed 4 doses/24 hrs	3 tablets every 6-8 hrs not to exceed 4 doses/24 hrs	3 tablets every 6-8 hrs not to exceed 4 doses/24 hrs

*Dosing is based primarily on weight, not age.